Few-Body Systems (EFB22) manuscript No.
(will be inserted by the editor)

C. Albertus

Semileptonic and nonleptonic decays of B_s mesons

arXiv:1403.2719v1 [hep-ph] 12 Mar 2014

Received: date / Accepted: date

Abstract We investigate the semileptonic and nonleptonic decays of B_s mesons. We work within the context of nonrelativistic constituent quark models. We calculate the different form factors that parameterize the hadron matrix elements.

Keywords Bottom · strange · semileptonic · nonleptonic

In this work we study the semileptonic decays of B_s mesons into D_s states and some of its nonleptonic decays. We work in the nonrelativistic constituent quark model scheme. Since the discovery of the B_s and $\bar{B}_s$ mesons, their lifetimes and decay modes have been in the aim of both theoretical and experimental collaboration and, for instance, are currently being measured in the LHCb experiment at CERN [1; 2; 3; 4; 5; 6; 7; 8; 9].

The description of the different J^P states involved is given in Ref. [10] and is omitted here for the shake of brevity. The semileptonic decays are governed by the V-A $b \to c$ current. The matrix elements are parametrized in terms of form factors as in Eq. (9) of Ref. [10] The calculation of the matrix elements and form factors can be found in full detail in Ref. [10], as well.

The semileptonic double differential decay width with respect to q^2 and the cosine, x_l, of the angle between the final meson momentum and the momentum of the final charged lepton, the latter measured in the lepton-neutrino center of mass frame (CMF), results to be, for a $\bar{B}_s$ at rest

$$\frac{d^2\Gamma}{dx_l dq^2} = \frac{G_F^2}{64m_{\bar{B}_s}^2}\frac{|V_{bc}|^2}{8\pi^3}\frac{\lambda^{1/2}(q^2, m_{\bar{B}_s}^2, m_{c\bar{s}}^2)}{2m_{\bar{B}_s}}\frac{q^2 - m_l^2}{q^2}\mathcal{H}_{\alpha\beta}(P_{\bar{B}_s}, P_{c\bar{s}})\mathcal{L}^{\alpha\beta}(p_l, p_\nu), \tag{1}$$

where G_F [11] is the Fermi constant, $\lambda(a,b,c) = (a+b-c)^2 - 4ab$, m_l is the mass of the charged lepton, $\mathcal{H}$ and $\mathcal{L}$ are the hadron and lepton tensors, and p_l and p_ν are the lepton four momenta. $P_{\bar{B}_s}$ and $P_{c\bar{s}}$ (with $c\bar{s} = D_s, D_s^*, D_{s2}^*$) are the meson four-momenta, $m_{\bar{B}_s}$ and $m_{c\bar{s}}$ their masses respectively, $P = P_{\bar{B}_s} + P_{c\bar{5}s}$, and $q = P_{\bar{B}_s} - P_{c\bar{s}}$. To calculate the contraction of the lepton and hadron tensors, that involve the form factors from Eq. (9) of [10], we shall follow the helicity formalism of Ref. [12]. This calculation is given in full detail in Sec. III E of Ref. [10]. Integrating Eq. 1 we obtain the total decay widths. Numerical values can be found in [10].

In this work we have also studied the nonleptonic $\bar{B}_s \to c\bar{s}M_F$ two-meson decays, where M_F is a pseudoscalar or vector meson. These decays correspond to a $b \to c$ transition at the quark level as well. These transitions are governed, neglecting penguin operators, by the effective Hamiltonian [13; 14]

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}}\left(V_{cb}\left[c_1(\mu)Q_1^{cb} + c_2(\mu)Q_2^{cb}\right] + H.c.\right), \tag{2}$$

C. Albertus
Departamento de Física Atómica, Molecular y Nuclear
Facultad de Ciencias. Universidad de Granada. Avenida de Fuentenueva S/N 37008 Granada Spain E-mail: albertus@ugr.es

Table 1 Branching fractions for the indicated decay channels, in percentage.

	BR in %	BR in %	
$\bar{B}_s \to D_s^+ e^- \bar{\nu}_e$	2.32	$\bar{B}_s \to D_s^+ \pi^-$	0.53
$\bar{B}_s \to D_s^{*+} e^- \bar{\nu}_e$	6.26	$\bar{B}_s \to D_s^+ \rho^-$	1.26
$\bar{B}_s \to D_s^+ \tau^- \bar{\nu}_\tau$	0.67	$\bar{B}_s \to D_s^+ K^-$	0.04
$\bar{B}_s \to D_s^{*+} \tau^- \bar{\nu}_\tau$	1.53	$\bar{B}_s \to D_s^+ K^{*-}$	0.08
$\bar{B}_s \to D_{s0}^{*+} \mu^- \bar{\nu}_\mu$	0.39	$\bar{B}_s \to D_{s0}^{*+} \pi^-$	0.10
$\bar{B}_s \to D_{s1}^{*+}(2460) \mu^- \bar{\nu}_\mu$	0.60	$\bar{B}_s \to D_{s0}^{*+} \rho^-$	0.27
$\bar{B}_s \to D_{s1}^{*+}(2536) \mu^- \bar{\nu}_\mu$	0.19	$\bar{B}_s \to D_{s0}^{*+} K^-$	0.009
$\bar{B}_s \to D_{s2}^{*+} \mu^- \bar{\nu}_\mu$	0.44	$\bar{B}_s \to D_{s0}^{*+} K^{*-}$	0.16

where $c_{1,2}$ are scale-dependent Wilson coefficients, and $Q_{1,2}$ are local four-quark operators given by

$$\begin{aligned} Q_1^{cb} =& \bar{\Psi}_c(0)\gamma_\mu(I-\gamma_5)\Psi_b(0)\left[V_{ud}^*\bar{\Psi}_d(0)\gamma^\mu(I-\gamma_5)\Psi_u(0) + V_{us}^*\bar{\Psi}_s(0)\gamma^\mu(I-\gamma_5)\Psi_u(0)\right. \\ & \left. + V_{cd}^*\bar{\Psi}_d(0)\gamma^\mu(I-\gamma_5)\Psi_c(0) + V_{cs}^*\bar{\Psi}_s(0)\gamma^\mu(I-\gamma_5)\Psi_c(0)\right] \\ Q_2^{cb} =& \bar{\Psi}_d(0)\gamma_\mu(I-\gamma_5)\Psi_b(0)\left[V_{ud}^*\bar{\Psi}_c(0)\gamma^\mu(I-\gamma_5)\Psi_u(0) + V_{cd}^*\bar{\Psi}_c(0)\gamma^\mu(I-\gamma_5)\Psi_c(0)\right] \\ & + \bar{\Psi}_s(0)\gamma_\mu(I-\gamma_5)\Psi_b(0)\left[V_{us}^*\bar{\Psi}_c(0)\gamma^\mu(I-\gamma_5)\Psi_u(0) + V_{cs}^*\bar{\Psi}_c(0)\gamma^\mu(I-\gamma_5)\Psi_c(0)\right], \end{aligned} \tag{3}$$

where V_{ij} are CKM matrix elements. We follow the factorization approximation, i. e., the hadron matrix elements of the effective Hamiltonian are evaluated as a product of quark-current matrix elements. One of these is that of the B_s transition to one of the final mesons, while the other one corresponds to a transition of the vacuum to the second final mesons, which is given by the corresponding meson decay constant. For $M_F = \pi, \rho, K$ or K^* the decay width is given by

$$\Gamma = \frac{G_F^2}{16\pi m_{B_s}^2}|V_{bc}|^2|V_F|^2\frac{\lambda^{1/2}(m_{B_s}^2, m_{c\bar{s}}^2, m_{M_F})^2}{2m_{B_s}}a_1^2\mathcal{H}_{\alpha\beta}(P_{B_s}, P_{c\bar{s}})\hat{\mathcal{H}}^{\alpha\beta}(P_F), \tag{4}$$

where $a_1 \approx 1.14$ is a Wilson coefficient and V_F is a CKM matrix elements which depends on the actual decay considered. As explained in Ref. [10], the contraction of the resulting tensors has been performed using the helicity formalism.

Some results for the semileptonic and nonleptonic decay widths can be found in Table 1. More results and comparison with previous calculations have been published in Ref. [10].

Acknowledgements The author thanks a contract granted by the CPAN project and support from Universidad de Granada and Junta de Andalucía, through the grant FQM-225.

References

1. R Aaij et al. *Phys.Rev.*, D87:112009, 2013.
2. R Aaij et al. *Phys.Rev.Lett.*, 2013.
3. Diego Guadagnoli and Gino Isidori. *eprint hep-ph://1302.3909*, 2013.
4. Steven Blusk. *eprint hep-ex://1212.4180*, 2012.
5. Alastair Dewhurst. *PoS*, BEAUTY2011:013, 2011.
6. Georges Aad et al. *JHEP*, 1212:072, 2012.
7. Gavril Giurgiu. *PoS*, FPCP2010:014, 2010.
8. A. Abulencia et al. *Phys.Rev.Lett.*, 96:191801, 2006.
9. F. Abe et al. *Phys.Rev.Lett.*, 74:4988, 1995.
10. C. Albertus. *e-print hep-ph://1401.1791*, 2014.
11. J. Beringer *et al. Phys. Rev. D*, 86:010001, Jul 2012.
12. Mikhail A. Ivanov, Juergen G. Korner, and Pietro Santorelli. *Phys.Rev.*, D71:094006, 2005.
13. D. Ebert, R.N. Faustov, and V.O. Galkin. *Phys.Rev.*, D75:074008, 2007.
14. M. Beneke, G. Buchalla, M. Neubert, and Christopher T. Sachrajda. *Phys.Rev.Lett.*, 83:1914–1917, 1999.